



**You have downloaded a document from  
RE-BUŚ  
repository of the University of Silesia in Katowice**

**Title:** On the random integral equation with advancing argument

**Author:** Henryk Gacki

**Citation style:** Gacki Henryk. (1985). On the random integral equation with advancing argument. "Annales Mathematicae Silesianae" (Nr 1 (1985), s. 47-50).



Uznanie autorstwa - Użycie niekomercyjne - Bez utworów zależnych Polska - Licencja ta zezwala na rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych oraz pod warunkiem zachowania go w oryginalnej postaci (nie tworzenia utworów zależnych).



UNIwersYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego

HENRYK GACKI

## ON THE RANDOM INTEGRAL EQUATION WITH ADVANCING ARGUMENT

**Abstract.** In this paper the existence and uniqueness of stochastic solutions of equation (1) are presented. Our theorem is an application of Banach fixed point theorem to the integral equation with advancing argument.

**Introduction.** Let  $(\Omega, \mathcal{B}, P)$  be a probability space. Denote by  $R^+ = [0, +\infty)$ ,  $R = (-\infty, +\infty)$ . We are mainly concerned with the existence and uniqueness of a random solution (defined below) of the random integral equation with advancing argument

$$(1) \quad x(\omega, t) = -y(\omega, t) + \int_0^{t+\delta(t)} K(\omega, u, x(\omega, u)) du,$$

where the kernel  $K$  is defined on  $\Omega \times R^+ \times R$ .

A mapping  $x: \Omega \times R^+ \rightarrow R$  is called a *random solution of equation (1)* if  $x(\omega, t)$  is a random variable for every  $t$  and if  $x$  satisfies equation (1) a.e. ( $P$ ). The mapping  $\bar{x}$  is the unique random solution of equation (1) if for every solution  $x$  of equation (1) and for every  $t \in R^+$ , the condition

$$\bar{x}(\omega, t) = x(\omega, t) \text{ a.e.}$$

holds.

Determined functional differential equations with advancing argument have been investigated by other authors (see [2], [3], [4]). The aim of this paper is to obtain the existence and uniqueness of the random solution of a random integral equation with advancing argument.

The main result of the paper is the following theorem.

**THEOREM.** *Suppose that the following assumptions are satisfied:*

---

*Received April 17, 1980.*

AMS (MOS) subject classification (1980). Primary 60H20.

1° For each  $(\omega, u) \in \Omega \times R^+$ ,  $K(\omega, u, \cdot)$  is a continuous function on  $R$ , and for each fixed  $v \in R$ ,  $K(\cdot, \cdot, v)$  is product-measurable on  $\Omega \times R^+$  and

$$\int_0^\infty \sup_v K(\omega, u, v) du < \infty.$$

2°  $\delta(t)$  is a nonnegative continuous function,  $\delta(0) = 0$ .

3° There exists a function  $a: \Omega \times R^+ \rightarrow R^+$  such that for each  $\omega \in \Omega$   $a(\omega, \cdot)$  is an integrable function and for each fixed  $u \in R^+$   $a(\cdot, u)$  is measurable on  $\Omega$ ; in addition the inequality

$$|K(\omega, u, v_1) - K(\omega, u, v_2)| \leq a(\omega, u)|v_1 - v_2|, \quad \omega \in \Omega, \quad u \in R^+, \quad v_1, v_2 \in R,$$

holds.

4° There exists  $0 < \Lambda < \frac{1}{e}$  such that,

$$|b(\omega, t + \delta(t)) - b(\omega, t)| < \Lambda, \quad \omega \in \Omega, \quad t \in R^+$$

where

$$b(\omega, t) = \int_0^t a(\omega, u) du.$$

5° For every  $\omega \in \Omega$ ,  $u \in R^+$  the inequality

$$|K(\omega, u, 0)| \leq ka(\omega, u) e^{eb(\omega, u)}$$

holds.

6° For each  $\omega \in \Omega$  the function  $y(\omega, \cdot)$  is continuous and for each  $t \in R^+$   $y(\cdot, t)$  is a random function; in addition, for fixed  $p$  the inequality

$$\sup_{t \geq 0} |y(\omega, t) e^{-b(\omega, t)}| \leq \frac{p}{2}$$

holds, where

$$p > \frac{2ke^{e\Lambda}}{e-2}.$$

Then equation (1) has the unique random solution.

Proof. Denote by  $\Phi$  the class of all mappings  $x: \Omega \times R^+ \rightarrow R$  such that for each  $\omega \in \Omega$   $x(\omega, \cdot) \in C(R^+, R)$ . For  $x \in \Phi$  and for fixed  $\omega \in \Omega$ , let

$$\|x\|_\omega = \sup_{t \geq 0} |x(\omega, t)| e^{-eb(\omega, t)}.$$

Let, for fixed  $\omega \in \Omega$ ,  $\Phi_\omega$ ,  $V_\omega$  denote subsets of  $C(R^+, R)$  such that  $\|x\|_\omega < \infty$  and  $\|x\|_\omega < p$ , respectively.

With the norm  $\|\cdot\|_\omega$ ,  $\Phi_\omega$  is a Banach space. It is easy to verify that  $V_\omega$  is a nonempty closed bounded subset of  $\Phi_\omega$ . Let  $E$  be the set of all mappings  $x \in \Phi$  such that, for each  $\omega \in \Omega$ ,  $x(\omega, \cdot) \in V_\omega$  and for each  $t \in R^+$   $x(\cdot, t)$  is a random variable. Then  $x \in E$  is product measurable on  $\Omega \times R^+$ . By 1°, the function

$$(\omega, t) \rightarrow \int_0^{t+\delta(t)} K(\omega, u, x(\omega, u)) du$$

is a random variable for each  $t \in R^+$  and for every  $\omega$  it is continuous in  $t$ . Define the operator  $T: \Phi \rightarrow \Phi$

$$(Tx)(\omega, t) = -y(\omega, t) + \int_0^{t+\delta(t)} K(\omega, u, x(\omega, u)) du.$$

The operator  $T$  has the following properties:

- (i)  $T(E) \subset E$ .
- (ii) For every  $x, z \in E$

$$\|Tx - Tz\|_\omega \leq L \|x - z\|_\omega,$$

where  $0 < L < 1$ .

Now, we shall prove (i). Let  $x \in E$ . Applying condition 3° we obtain

$$|K(\omega, u, x(\omega, u)) - K(\omega, u, 0)| \leq a(\omega, u) e^{eb(\omega, u)} \|x\|_\omega.$$

In virtue of 4° and 5° we have

$$\begin{aligned} |(Tx)(\omega, t)| &< |y(\omega, t)| + \int_0^{t+\delta(t)} |K(\omega, u, x(\omega, u))| du \leq \\ &\leq |y(\omega, t)| + \int_0^{t+\delta(t)} e |a(\omega, u)| e^{eb(\omega, u)} du \frac{\|x\|_\omega}{e} + \frac{k}{e} \int_0^{t+\delta(t)} e |a(\omega, u)| e^{eb(\omega, u)} du \leq \\ &\leq |y(\omega, t)| + \left( \frac{\|x\|_\omega}{e} + \frac{k}{e} \right) e^{eb(\omega, t+\delta(t))}. \end{aligned}$$

By conditions 4° and 5° it follows that

$$\|Tx\|_\omega \leq \|y\|_\omega + \left( \frac{\|x\|_\omega}{e} + \frac{k}{e} \right) e^{eA} \leq \frac{p}{2} + \left( \frac{p}{e} + \frac{k}{e} \right) e^{eA} \leq p.$$

Now, we shall prove (ii). Let  $x, z \in E$ . Applying the Lipschitz conditions 2° and 3° we have:

$$\begin{aligned} |(Tx)(\omega, t) - (Tz)(\omega, t)| &\leq \int_0^{t+\delta(t)} |a(\omega, u)| e^{eb(\omega, u)} du \|x - z\|_\omega \leq \\ &\leq \frac{\|x - z\|_\omega}{e} e^{eb(\omega, t+\delta(t))}. \end{aligned}$$

This inequality implies that

$$\|Tx - Tz\|_\omega \leq \frac{e^{eA}}{e} \|x - z\|_\omega.$$

Now, define

$$L = \frac{e^{eA}}{e} < 1.$$

Then defining recursively the sequence:

$$\begin{aligned} x_0(\omega, t) &= -y(\omega, t), \\ &\dots\dots\dots, \\ x_{n+1}(\omega, t) &= -y(\omega, t) + \int_0^{t+\delta(t)} K(\omega, u, x_n(\omega, u)) du, \end{aligned}$$

we see that all mappings  $x_n$  are in  $E$  and, for each  $\omega$ ,  $x_n(\omega, \cdot)$  is a Cauchy sequence in the Banach space  $(\Phi_\omega, \|\cdot\|_\omega)$ . Hence, there exists an  $x$  in  $E$  such that for each  $\omega \in \Omega$

$$x(\omega, \cdot) = \lim_{n \rightarrow \infty} x_n(\omega, \cdot) \text{ in } (\Phi_\omega, \|\cdot\|_\omega).$$

Finally, we shall show that  $x$  is a fixed point of  $T$ . Indeed, for  $n$  large enough, we have

$$\|x_n - Tx\|_\omega \leq \|Tx_{n-1} - Tx\|_\omega \leq L \|x_{n-1} - x\|_\omega$$

and

$$\|Tx - x\|_\omega \leq \|Tx - x_n\|_\omega + \|x_n - x\|_\omega,$$

hence  $\|Tx - x\|_\omega = 0$  i.e.

$$\sup_{t \geq 0} e^{-eb(\omega, t)} |(Tx)(\omega, t) - x(\omega, t)| = 0 \text{ for every } \omega \in \Omega.$$

Consequently,

$$(Tx)(\omega, t) - x(\omega, t) = 0.$$

Thus  $x$  is the unique solution of (1). This completes the proof.

*Acknowledgment.* The author is indebted to Prof. J. Błaż for encouragement and several useful remarks.

#### REFERENCES

- [1] A. T. BHARUCHA-REID, A. MUKHEREJEA and N. A. TSERPES, *On the existence of random solutions of some random integral equations*, Rev. Roum. Math. Pures et Appl. 22 (1977).
- [2] A. BIELECKI, J. BŁAŻ, *Über eine Verallgemeinerung der Nicoletti-Aufgabe für Funktional-Differentialgleichungen mit voreilendem Argument*, Monatsh. Math. 88 (1979), 287—291.
- [3] J. BŁAŻ, *Über die Nicoletti-Aufgabe für Funktional-Differentialgleichungen mit voreilendem Argument*, Arch. Math. 27 (1976), 530—543.
- [4] J. BŁAŻ, W. WALTER, *Über Funktional-Differentialgleichungen mit voreilendem Argument*, Monatsh. Math. 82 (1976), 1—16.